

Magnetic monopole loops supported by a meron pair as the quark confiner

Kei-Ichi Kondo*

Department of Physics, Graduate School of Science, Chiba University, Chiba 263-8522, Japan

E-mail: kondok@faculty.chiba-u.jp

We give a definition of gauge-invariant magnetic monopoles in Yang-Mills theory without using the Abelian projection due to 't Hooft. They automatically appear from the Wilson loop operator. This is shown by rewriting the Wilson loop operator using a non-Abelian Stokes theorem. The magnetic monopole defined in this way is a topological object of co-dimension 3, i.e., a loop in four-dimensions. We show that such magnetic loops indeed exist in four-dimensional Yang-Mills theory. In fact, we give an analytical solution representing circular magnetic monopole loops joining a pair of merons in the four-dimensional Euclidean $SU(2)$ Yang-Mills theory. This is achieved by solving the differential equation for the adjoint color (magnetic monopole) field in the two-meron background field within the recently developed reformulation of the Yang-Mills theory. Our analytical solution corresponds to the numerical solution found by Montero and Negele on a lattice. This result strongly suggests that a meron pair is the most relevant quark confiner in the original Yang-Mills theory, as Callan, Dashen and Gross suggested long ago.

*8th Conference Quark Confinement and the Hadron Spectrum
September 1-6 2008
Mainz, Germany*


*This work is financially supported by Grant-in-Aid for Scientific Research (C) 18540251 from Japan Society for the Promotion of Science (JSPS).

1. Wilson loop and magnetic monopole

For a closed loop C , the Wilson loop operator for $SU(2)$ Yang-Mills connection is defined by

$$W_C[\mathbf{A}] := \text{tr} [\mathcal{P} \exp \{ig \oint_C dx^\mu \mathbf{A}_\mu(x)\}] / \text{tr}(\mathbf{1}), \quad \mathbf{A}_\mu(x) = \mathbf{A}_\mu^A(x) \sigma^A / 2. \quad (1.1)$$

The path-ordering $\mathcal{P}$ is removed by using the Diakonov-Petrov version [1] of a non-Abelian Stokes theorem for the Wilson loop operator: in the J representation of $SU(2)$ ($J = 1/2, 1, 3/2, 2, \dots$)

$$\begin{aligned} W_C[\mathbf{A}] &:= \int d\mu_\Sigma(U) \exp \left\{ iJg \int_{\Sigma: \partial\Sigma=C} dS^{\mu\nu} f_{\mu\nu} \right\}, \text{ no path-ordering} \\ f_{\mu\nu}(x) &:= \partial_\mu [\mathbf{A}_\nu^A(x) \mathbf{n}^A(x)] - \partial_\nu [\mathbf{A}_\mu^A(x) \mathbf{n}^A(x)] - g^{-1} \varepsilon^{ABC} \mathbf{n}^A(x) \partial_\mu \mathbf{n}^B(x) \partial_\nu \mathbf{n}^C(x), \\ \mathbf{n}^A(x) \sigma^A &:= U^\dagger(x) \sigma^3 U(x), \quad U(x) \in SU(2) \quad (A, B, C \in \{1, 2, 3\}), \end{aligned} \quad (1.2)$$


where $d\mu_\Sigma(U)$ is the product measure of an invariant measure on $SU(2)/U(1)$ over Σ :

$$d\mu_\Sigma(U) := \prod_{x \in \Sigma} d\mu(U(x)), \quad d\mu(U(x)) = \frac{2J+1}{4\pi} \delta(\mathbf{n}^A(x) \mathbf{n}^A(x) - 1) d^3 \mathbf{n}(x), \quad (1.3)$$

where we have introduced a unit vector field $\mathbf{n}(x)$.

The geometric and topological meaning of the Wilson loop operator was given in [2]:

$$W_C[\mathcal{A}] = \int d\mu_\Sigma(U) \exp \{iJg(\Xi_\Sigma, k) + iJg(N_\Sigma, j)\}, \quad C = \partial\Sigma \quad (1.4)$$

$$k := \delta^* f = *df, \quad \Xi_\Sigma := \delta^* \Theta_\Sigma \Delta^{-1} \leftarrow \text{(D-3)-forms} \quad (1.5)$$

$$j := \delta f, \quad N_\Sigma := \delta \Theta_\Sigma \Delta^{-1} \leftarrow \text{1-forms (D-indep.)} \quad (1.6)$$

$$\Theta_\Sigma^{\mu\nu}(x) = \int_\Sigma d^2 S^{\mu\nu}(x(\sigma)) \delta^D(x - x(\sigma)), \quad (1.7)$$

where k and j are gauge invariant and conserved currents, $\delta k = 0 = \delta j$. Thus, **we do not need to use the Abelian projection proposed by 't Hooft [3] to define magnetic monopoles in Yang-Mills theory! The Wilson loop operator knows the (gauge-invariant) magnetic monopole!**

Then the magnetic monopole is a topological object of co-dimension 3. In D dimensions,

D=3: 0-dimensional point defect $\rightarrow$ magnetic monopole of Wu-Yang type

D=4: 1-dimensional line defect $\rightarrow$ magnetic monopole loop (closed loop)

For $D = 3$,

$$k(x) = \frac{1}{2} \varepsilon^{jk\ell} \partial_\ell f_{jk}(x) = \rho_m(x) \quad (1.8)$$

denotes the magnetic charge density at x , and


$$\Xi_\Sigma(x) = \Omega_\Sigma(x) / (4\pi) \quad (1.9)$$

agrees with the (normalized) solid angle at the point x subtended by the surface Σ bounding the Wilson loop C . Then the magnetic part $W_{\mathcal{A}}^m$ is written as

$$W_{\mathcal{A}}^m := \exp \{iJg(\Xi_\Sigma, k)\} = \exp \left\{ iJg \int d^3 x \rho_m(x) \frac{\Omega_\Sigma(x)}{4\pi} \right\}. \quad (1.10)$$

The magnetic charge q_m obeys the Dirac-like quantization condition:

$$q_m := \int d^3 x \rho_m(x) = 4\pi g^{-1} n \quad (n \in \mathbb{Z}). \quad (1.11)$$



The proof follows from a fact that the non-Abelian Stokes theorem does not depend on the surface Σ chosen for spanning the surface bounded by the loop C . See [2].

For an ensemble of point-like magnetic charges: $k(x) = \sum_{a=1}^n q_m^a \delta^{(3)}(x - z_a)$, we have

$$W_{\mathcal{A}}^m = \exp \left\{ iJ \frac{g}{4\pi} \sum_{a=1}^n q_m^a \Omega_{\Sigma}(z_a) \right\} = \exp \left\{ iJ \sum_{a=1}^n n_a \Omega_{\Sigma}(z_a) \right\}, \quad n_a \in \mathbb{Z}. \quad (1.12)$$

The magnetic monopoles in the neighborhood of the Wilson surface Σ ($\Omega_{\Sigma}(z_a) = \pm 2\pi$) contribute to the Wilson loop

$$W_{\mathcal{A}}^m = \prod_{a=1}^n \exp(\pm i2\pi J n_a) = \begin{cases} \prod_{a=1}^n (-1)^{n_a} & (J = 1/2, 3/2, \dots) \\ = 1 & (J = 1, 2, \dots) \end{cases}. \quad (1.13)$$

This enables us to explain the N -ality dependence of the asymptotic string tension. See, [4].

For $D = 4$, $\Omega_{\Sigma}^{\mu}(x)$ is the $D = 4$ solid angle and the magnetic part reads

$$W_{\mathcal{A}}^m = \exp \left\{ iJg \int d^4x \Omega_{\Sigma}^{\mu}(x) k^{\mu}(x) \right\}. \quad (1.14)$$

Suppose the existence of an ensemble of magnetic monopole loops C'_a in $D = 4$ Euclidean space, $k^{\mu}(x) = \sum_{a=1}^n q_m^a \oint_{C'_a} dy_a^{\mu} \delta^{(4)}(x - x_a)$, $q_m^a = 4\pi g^{-1} n_a$. Then the Wilson loop operator reads

$$W_{\mathcal{A}}^m = \exp \left\{ iJg \sum_{a=1}^n q_m^a L(\Sigma, C'_a) \right\} = \exp \left\{ 4\pi Ji \sum_{a=1}^n n_a L(\Sigma, C'_a) \right\}, \quad n_a \in \mathbb{Z}, \quad (1.15)$$

where $L(\Sigma, C')$ is the linking number between the surface Σ and the curve C' :


$$L(\Sigma, C') := \oint_{C'} dy^{\mu}(\tau) \Xi_{\Sigma}^{\mu}(y(\tau)). \quad (1.16)$$

Here the curve C' is identified with the trajectory k of a magnetic monopole and the surface Σ with the world sheet of a hadron (meson) string for a quark-antiquark pair.

The Wilson loop operator is a probe of the gauge-invariant magnetic monopole defined in our formulation. Thus, calculating the Wilson loop average reduces to the summation over the magnetic monopole charge ($D=3$) or current ($D=4$) with a geometric factor, the solid angle ($D=3$) or linking number ($D=4$).

2. Main results (Magnetic loops indeed exist in YM_4)

We can show that **the gauge-invariant magnetic loop (assumed in the above) indeed exists in $SU(2)$ Yang-Mills theory in $D = 4$ Euclidean space**: we give a first* (exact) analytical solution representing **circular magnetic monopole loops joining two merons** [5].¹



Our method reproduces also the previous results based on MAG (MCG) and LAG:

- (i) The magnetic straight line can be obtained in the one-instanton or one-meron background. [6, 7]
- (ii) The magnetic closed loop can NOT be obtained in the one-instanton background. [8, 9]

¹There is an exception: Bruckmann & Hansen, hep-th/0305012, Ann.Phys.**308**, 201 (2003). However, it has $Q_P = \infty$

3. Reformulating Yang-Mills theory in terms of new variables

SU(2) Yang-Mills theory $\longleftrightarrow$ A reformulated Yang-Mills theory
 written in terms of $\mathbf{A}_\mu^A(x)$ ($A = 1, 2, 3$) $\longleftrightarrow$ written in terms of new variables:
 $\mathbf{A}_\mu^A(x)$ ($A = 1, 2, 3$) $\longleftrightarrow$ change of variables $\mathbf{n}^A(x), c_\mu(x), \mathbf{X}_\mu^A(x)$ ($A = 1, 2, 3$)
 We introduce a “color field” $\mathbf{n}(x)$ of unit length with three components

$$\mathbf{n}(x) = (n_1(x), n_2(x), n_3(x)), \quad \mathbf{n}(x) \cdot \mathbf{n}(x) = n_A(x)n_A(x) = 1 \quad (3.1)$$

The color field $\mathbf{n}(x)$ is identified with $\mathbf{n}(x)$ in (1.2). New variables $\mathbf{n}^A(x), c_\mu(x), \mathbf{X}_\mu^A(x)$ should be given as functionals of the original $\mathbf{A}_\mu^A(x)$. The off-shell Cho-Faddeev-Niemi-Shabanov decomposition [10] is reinterpreted as change of variables from $\mathbf{A}_\mu^A(x)$ to $\mathbf{n}^A(x), c_\mu(x), \mathbf{X}_\mu^A(x)$ via the reduction of an enlarged gauge symmetry. See [11, 12]. Expected role of the color field: 1) The color field $\mathbf{n}(x)$ plays the role of recovering color symmetry which will be lost in the conventional approach, e.g., in the MA gauge. 2) The color field $\mathbf{n}(x)$ carries topological defects responsible for non-perturbative phenomena, e.g., quark confinement.

4. Bridge between $\mathbf{A}_\mu(x)$ and $\mathbf{n}(x)$

For a given Yang-Mills field $\mathbf{A}_\mu(x)$, the color field $\mathbf{n}(x)$ is obtained by solving the reduction differential equation (RDE): [12]

$$\mathbf{n}(x) \times D_\mu[\mathbf{A}]D_\mu[\mathbf{A}]\mathbf{n}(x) = \mathbf{0}. \quad (4.1)$$

For a given SU(2) Yang-Mills field $\mathbf{A}_\mu(x) = \mathbf{A}_\mu^A(x) \frac{\sigma_A}{2}$, look for unit vector fields $\mathbf{n}(x)$ such that $-D_\mu[\mathbf{A}]D_\mu[\mathbf{A}]\mathbf{n}(x)$ is proportional to $\mathbf{n}(x)$: an eigenvalue-like form:


$$-D_\mu[\mathbf{A}]D_\mu[\mathbf{A}]\mathbf{n}(x) = \lambda(x)\mathbf{n}(x) \quad (\lambda(x) \geq 0). \quad (4.2)$$

The solution is not unique. We choose the solution giving the smallest value of the reduction functional F_{rc} which agrees with the integral of the scalar function $\lambda(x)$ over $\mathbb{R}^D \mathbb{A}F$

$$\begin{aligned}
 F_{\text{rc}} &= \int d^D x \frac{1}{2} (D_\mu[\mathbf{A}]\mathbf{n}(x)) \cdot (D_\mu[\mathbf{A}]\mathbf{n}(x)) = \int d^D x \frac{1}{2} \mathbf{n}(x) \cdot (-D_\mu[\mathbf{A}]D_\mu[\mathbf{A}]\mathbf{n}(x)) \\
 \implies F_{\text{rc}}^* &= \int d^D x \frac{1}{2} \mathbf{n}(x) \cdot \lambda(x)\mathbf{n}(x) = \int d^D x \frac{1}{2} \lambda(x).
 \end{aligned} \quad (4.3)$$


5. Conclusion and discussion

For given one-instanton and two-meron background $\mathbf{A}_\mu(x)$, we have solved the RDE for the color field $\mathbf{n}(x)$ [12]. In the four-dimensional Euclidean SU(2) Yang-Mills theory, we have given a first analytical solution representing circular magnetic monopole loops k_μ which go through a pair of merons (with a unit topological charge) with non-trivial linking with the Wilson surface Σ .




This is achieved by solving the reduction differential equation for the adjoint color (magnetic monopole) field in the two-meron background field using the recently developed reformulation of the Yang-Mills theory [11, 12] and a non-Abelian Stokes theorem [2].

Our analytical solution corresponds to a numerical solution found on a lattice by Montero and Negele [13].



We have not yet obtained the analytic solution representing magnetic loops connecting 2-instantons, which were found in the numerical way by Reinhardt & Tok [7].



Thus we are lead to a conjecture: A meron pair is the most relevant quark confiner in the original Yang-Mills theory, as Callan, Dashen and Gross suggested long ago [14]. This means a duality relation:

$$\text{dual Yang-Mills: magnetic monopole loops} \iff \text{original Yang-Mills: merons}$$

References

- [1] D. Diakonov & V. Petrov, Phys. Lett. **B224**, 131 (1989); hep-th/9606104.
- [2] K.-I. Kondo, arXiv:0801.1274, Phys.Rev.D**77**, 085029 (2008). K.-I. Kondo, hep-th/0009152.
- [3] G. 't Hooft, Nucl. Phys. **B190**, 455 (1981).
- [4] K.-I. Kondo, arXiv:0802.3829, J.Phys.G**35**, 085001 (2008).
- [5] K.-I. Kondo, N. Fukui, A. Shibata & T. Shinohara, arXiv:0806.3913, Phys. Rev. D**78**, 065033 (2008).
- [6] M. Chernodub & F. Gubarev, hep-th/9506026, JETP Lett. **62**, 100 (1995).
- [7] H. Reinhardt & T. Tok, hep-th/0011068, Phys.Lett.B**505**, 131 (2001).
- [8] R.C. Brower, K.N. Orginos & C-I. Tan, hep-th/9610101, Phys.Rev.D **55**, 6313 (1997).
- [9] F. Bruckmann, T. Heinzl, T. Vekua & A. Wipf, hep-th/0007119, Nucl.Phys.B**593**, 545 (2001).
- [10] Y. M. Cho, Phys. Rev. D **21**, 1080 (1980); **23**, 2415 (1981); L. D. Faddeev and A. J. Niemi, Phys. Rev. Lett. **82**, 1624 (1999); Nucl. Phys. B**776**, 38 (2007); S.V. Shabanov, Phys. Lett. B **458**, 322 (1999); Phys. Lett. B**463**, 263 (1999).
- [11] K.-I. Kondo, T. Murakami & T. Shinohara, hep-th/0504107, Prog. Theor. Phys. **115**, 201 (2006).
- [12] K.-I. Kondo, T. Shinohara & T. Murakami, arXiv:0803.1076, Prog. Theor. Phys. **120**,1 (2008).
- [13] A. Montero & J.W. Negele, hep-lat/0202023, Phys.Lett.B**533**, 322 (2002).
- [14] C.G. Callan, R. Dashen and D.J. Gross, Phys. Rev. D **17**, 2717 (1978).